Peninsula Master Gardener Small Fruit Class

Anne Geyer 2006 Hanover Master Gardener Founder of Agriberry Farm Hanover VA Email: <u>afgeyer@yahoo.com</u> Cell 804-314-7517



Real Inspire and instruct on tidewater strawberries, brambles, blueberries, grapes and minor fruits

Reprovide a resource list for expanding further knowledge

KEY RESOURCE : THE VIRGINIA FRUIT PAGE





The Mid-Atlantic Regional Fruit Loop:

The Virginia Fruit Page

Extension, Research and Teaching in Fruit Crops

Fruit Production in the mid-Atlantic region has a long history, dating to Colonial times in Virginia with pomological and viticultural efforts by founders such as Thomas. Jefferson and George Washington. Today, Virginia ranks approximately sixth in the nation for apple production (about 11,029 acres). There are about 1,770 acres of peach, 4,370 acres of grape, and about 425 acres of blueberry, 380 acres of caneberries and 290 acres of strawberry (USDA 2012 census).

For more detailed information, please click on the desired crop:



Here is a simple list of links to resources available in this fruit home page:

- Virginia Tech Tree Fruit Extension and Outreach Web Site
- Virginia Tech Tree Fruit Extension and Outreach in Facebook
- Home Fruit Production
- Horticultural Practices
- Fruit IPM
- Honey Bees and Pollination
- Weather Information
 - National Climate Assessment (May 2014 report)
- Soil Testing
- Mid-Atlantic Land-Grant University Fruit Faculty
- Research Units
- Other Sources of Fruit Information
- New! Interested in Continuing Education?

MOST IMPORTANT RESOURCE

Fruit provides <u>perennial</u> opportunities for vertical development in the garden



Reasons to Promote Fruit Culture

Ramily's quality of life & health
Locally produced food and beverages
Value as landscape elements
Outdoor recreation
Preserve heirloom varieties
Fruit industry sustainability
Social interaction including CSA's

Incredible Edible Berries



Require minimum space for amount of fruit produced

Rever house of nutrients

Realize array of healthy textures, colors and flavors

Strawberries

Broad culture options
 Regular matted row planting ideal for home
 Straw mulch
 TIP : remove first flowers post planting



Raspberries

Red Raspberries
Purples Raspberries
Black Raspberries
Fall Raspberries





Purple Raspberries

Cornell Univ Cross between Black Raspberry and Red Raspberry in the 1970's - This is the variety ROYALTY



Black Raspberries

Black Raspberries have much higher health benefits than blueberries but much lower yields





Blackberries

-Acclimated to VA -Great new **Thornless Varieties** -Options for 4 season hands on management ??? -New varieties for Fall fruiting ©





Blueberries



pH is a vital factor here
Mulch is critical
Plant a 1 gal nursery plant







Grapes



 For Tidewater, try Muscadines or limited plantings
 Prepare for Pest issues





Currants, Gooseberries & Elderberries









Planting : Location Location Location



Consider IPM : 3 M's
Minimize
Monitor
Manage





Optimize your site when possible



Well Drained Soil is KEY And proper pH is too
















Weed management







Structure and support is key! 2400 posts...18 miles of wire You can be innovative ③







Caneberry Pruning in Feb, May, & September...



Ready for "Fall" Harvest Pick from May 12 - December 1









Common Pests

Pest" has a very broad scope in Berry Crops! PEST = any influence that can affect the plants life cycle

- people, deer, birds, sun, frost, rain, hail, wind, drought, dust, corn bin dryer...flood
- Apply authorized, up to date "Crop Protectants" (not Pesticides)

PRIMARY PEST MANAGEMENT CONSIDERATIONS

Correct identification of insects and diseases
 Determine if a need for control exists and proper timing of treatments

Sanitation

- The destruction of places that harbor insects and diseases plays a large part in the control program.
 Collect and burn debris.
- Remove and destroy all dropped fruit.
- Rake and burn apple and cherry leaves.
- Scrape loose bark from trunks, crotches, and main limbs of apple trees.
- Reverse out and destroy all dead or diseased
- Conditions that encourage mice should also be eliminated.

Hail Damage












Verticilium Wilt







Up to 66 known caneberry viruses...



Leaf Curl Virus ?



Fall Lady Beetles



SWD Spotted Wing Drosophila


New pest and now in most of US, including Virginia. Minimize with good sanitation Monitor with vinegar trap Manage with rotating pyrethrum insecticides







3-4 days after egg laid



Skin wrinkles and fruit becomes juicy (dark scarring could happen at 1-2 days)

After 5 days



Dark scarring





The Virginia Fruit Page

Extension, Research and Teaching in Fruit Crops Mid-Atlantic Regional Fruit Loop

http://www.virginiafruit.ento.vt.edu/









2005...NBCI vision to triple berry consumption by 2020?

- Without land or wealth could I be the first US berry farm started with a Community Supported Agriculture "CSA" model?
- Could I fund a first worker training program around summer raspberries and blackberries?
- Aware of Richmond's emerging locavore market AND "Free resources" for new businesses ...SBA, SCORE



AGRIBERRY MISSION:

TO ESTABLISH LOCAL CONSUMER SUPPORTED SUMMER BERRY FIELDS AND BE A PREMIER AGRICULTURAL TRAINING FACILITY FOR MOTIVATED FIRST TIME WORKERS.

Geyer Backyard Berries





20 Week Agriberry CSA









Young people involved











Small fruit growing & marketing is what its all about for the Agriberry family farm!



<complex-block>



anne@agriberry.com

cell 804-314-7517

www.agriberry.com